

## REPORT DOCUMENTATION PAGE

Public reporting burden for this collection of information is estimated to average 1 hour per response, including gathering and maintaining the data needed, and completing and reviewing the collection of information. Send collection of information, including suggestions for reducing this burden, to Washington Headquarters Service, Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paper

yes,  
this  
non

|  |  |  |  |  |  |
|--|--|--|--|--|--|
| 1. AGENCY USE ONLY (Leave blank)   |  | 2. REPORT DATE                           |  | 3. REPORT TYPE AND DATES COVERED               |  |
|  |  |  |  | Final 01 Jul 93 to 30 Jun 97                   |  |
| 4. TITLE AND SUBTITLE  |  |  |  | 5. FUNDING NUMBERS                             |  |
| (AASERT-92) ULTRA PRECISE LASER SPECTROSCOPY   |  |  |  | 61103D<br>3484/TS                              |  |
| 6. AUTHOR(S)   |  |  |  |  |  |
| Professor Kleppner   |  |  |  |  |  |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)   |  |  |  | 8. PERFORMING ORGANIZATION REPORT NUMBER       |  |
| Massachusetts Institute of Technology<br>77 Massachusetts Avenue<br>Cambridge MA 02139-4307  |  |  |  |  |  |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)  |  |  |  | 10. SPONSORING/MONITORING AGENCY REPORT NUMBER |  |
| AFOSR/NE<br>110 Duncan Avenue Rm B115<br>Bolling AFB DC 20332-8050   |  |  |  | F49620-93-1-0395                               |  |
| 11. SUPPLEMENTARY NOTES  |  |  |  |  |  |
|  |  |  |  |  |  |
| 12a. DISTRIBUTION / AVAILABILITY STATEMENT   |  |  |  | 12b. DISTRIBUTION CODE                         |  |
| APPROVAL FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED  |  |  |  |  |  |
| 13. ABSTRACT (Maximum 200 words)   |  |  |  |  |  |
| <p>Research on this AASERT grant was focussed on the optical spectroscopy of trapped cold hydrogen for use as a spectroscopic tool and as a means of studying the gas. The grant supported research by two graduate students. During the grant an optical signal on the 1S - 2S transition in cold trapped hydrogen was observed, opening the way to a new form of ultra high resolution spectroscopy. The optical signal will also provide a valuable tool for manipulating a Bose condensate in hydrogen. The condensate can absorb two photons form a single running wave. The recoil is enough to sweep the atoms out of the trap, creating a new type of atom laser beam.</p> |  |  |  |  |  |
| 14. 19980115 215   |  |  |  | 15. NUMBER OF PAGES                            |  |
|  |  |  |  |  |  |
|  |  |  |  | 16. PRICE CODE                                 |  |
|  |  |  |  |  |  |
| 17. SECURITY CLASSIFICATION OF REPORT  |  | 18. SECURITY CLASSIFICATION OF THIS PAGE |  | 19. SECURITY CLASSIFICATION OF ABSTRACT        |  |
| UNCLASSIFIED   |  | UNCLASSIFIED                             |  | UNCLASSIFIED                                   |  |
|  |  |  |  | 20. LIMITATION OF ABSTRACT                     |  |
|  |  |  |  | UL   |  |

**MASSACHUSETTS INSTITUTE OF TECHNOLOGY  
DEPARTMENT OF PHYSICS**

**DANIEL KLEPPNER**  
Lester Wolfe Professor of Physics  
Associate Director, Research Laboratory of Electronics

address: MIT  
room 26-237  
Cambridge, MA 02139

phone: 617/253-6811  
fax: 617/253-4876  
dk@amo.mit.edu

November 7, 1997

Dr. Ralph Kelley  
Air Force Office of Scientific Research  
Building 410  
Bolling Air Force Base  
Washington, DC 20332

Re: AASERT Grant F49620-93-1-0395, Final report

Dear Dr. Kelley,

This is the final report for the above AASERT grant.


The research under this grant is focussed on the optical spectroscopy of trapped cold hydrogen for use as a spectroscopic tool and as a means of studying the gas. The grant supported research by two outstanding graduate students, Dale Fried and Tom Killian. Both of these will prepare their theses in the near future.

During the period of this grant we succeeded in observing an optical signal on the 1S - 2S transition in cold trapped hydrogen, opening the way to a new form of ultra high resolution spectroscopy. The optical signal will also provide a valuable tool for manipulating a Bose condensate in hydrogen. The condensate can absorb two photons from a single running wave. The recoil is enough to sweep the atoms out of the trap, creating a new type of atom laser beam.

The final stages of this research involved a major redesign of our trapping cell, allowing us to implement rf cooling of the trapped hydrogen. The new trap is currently being brought into production.

A list of publications supported by this work is attached.

Yours sincerely,



Daniel Kleppner  
Lester Wolfe Professor of Physics

Doppler-Free Spectroscopy of Trapped Atomic Hydrogen, Thomas C. Killian, Dale G. Fried, Claudio L. Cesar, Adam D. Polcyn, Thomas J. Greytak and Daniel Kleppner, Atomic Physics 15, H.B. Van Linden Van Den Heuvell, J.T.M Walraven, M.W. Reynolds (World Scientific, Singapore, 1997), p. 158. ICAP 96.

Two-photon spectroscopy of trapped atomic hydrogen, Claudio L. Cesar, Dale G. Fried, Thomas C. Killian, Adam D. Polcyn, Jon C. Sandberg, Ite A. Yu, Thomas J. Greytak, Daniel Kleppner, and John Doyle, Phys. Rev. Lett. 77, 255 (1996)

Two-Photon Spectroscopy of Trapped Atomic Hydrogen, C.L. Cesar, D.G. Fried, T.C. Killian, A.D. Polcyn, J.C. Sandberg, J.M. Doyle, I.A. Yu, T.J. Greytak and D. Kleppner, Proceedings of the Fifth Symposium on Frequency Standards and Metrology, J.C. Bergquist, ed. (World Scientific, Singapore) 1996, p 365.